

! SAFETY ALERT !

HYDROXYCUT RECALL

IF YOU ARE USING HYDROXYCUT, STOP DOING SO IMMEDIATELY.

On 1 May 2009, the U.S. Food and Drug Administration (FDA) published a consumer warning advising consumers to ***immediately stop using*** Hydroxycut products. These products have been linked to at least 23 cases of serious liver injuries, including damage requiring liver transplant and one confirmed death.

If you have used Hydroxycut, the early signs of liver injury include loss of appetite, nausea, fatigue or weakness. Symptoms can progress to vomiting, brown urine, light-colored stools, and/or yellowing of the skin or whites of eyes (jaundice).

Other problems associated with Hydroxycut

products include seizures, heart problems, and muscle damage. The symptoms can occur at any dose and at any time.

Products with similar ingredients may also pose a risk of causing harmful health effects and should be avoided.

If you have any symptoms that could be associated with these or other dietary supplements, consult a physician or other health care professional. Report all adverse effects you've experienced to the FDA by going to <https://www.accessdata.fda.gov/scripts/medwatch/medwatch-online.htm> and send an email to supplements@usuhs.edu.

RECALLED PRODUCTS INCLUDE:

Hydroxycut Regular Rapid Release Caplets
Hydroxycut Caffeine-Free Rapid Release Caplets
Hydroxycut Hardcore Liquid Caplets
Hydroxycut Max Liquid Caplets
Hydroxycut Regular Drink Packets
Hydroxycut Caffeine-Free Drink Packets
Hydroxycut Hardcore Drink Packets (Ignition Stix)
Hydroxycut Max Drink Packets
Hydroxycut Liquid Shots
Hydroxycut Hardcore RTDs (Ready-to-Drink)
Hydroxycut Max Aqua Shed
Hydroxycut 24
Hydroxycut Carb Control
Hydroxycut Natural



For more information:

<http://www.fda.gov/consumer/updates/hydroxycut050109.pdf>

<http://www.iovate.com/supplements@usuhs.edu>